


WHAT IS CLAIMED IS

1. An agent for the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis, which comprises a compound having a Rho kinase inhibitory activity.

5




2. The agent for the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis of Claim 1, wherein the compound having a Rho kinase inhibitory activity is an amide compound of the following formula (I)



10


wherein

Ra is a group of the formula



in the formulas (a) and (b),

15 R is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring, or a group of the formula



wherein R⁶ is hydrogen, alkyl or formula: -NR⁸R⁹


wherein R⁸ and R⁹ are the same or different and each is hydrogen, alkyl, aralkyl or phenyl, R⁷ is hydrogen, alkyl, aralkyl, phenyl, nitro or cyano, or R⁶ and R⁷ in combination show a group forming a heterocycle optionally having, in the ring, oxygen atom, sulfur atom or optionally substituted nitrogen atom,

R¹ is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring, or R and R¹ in combination form, together with the adjacent nitrogen atom, a group forming a heterocycle optionally having, in the ring, oxygen atom, sulfur atom or optionally substituted nitrogen atom,

R² is hydrogen or alkyl,

R³ and R⁴ are the same or different and each is hydrogen, alkyl, aralkyl, halogen, nitro, amino, alkylamino, acylamino, hydroxy, alkoxy, aralkyloxy, cyano, acyl, mercapto, alkylthio, aralkylthio, carboxy, alkoxycarbonyl, carbamoyl, alkylcarbamoyl or azide, and

A is a group of the formula







wherein R¹⁰ and R¹¹ are the same or different and each is hydrogen, alkyl, haloalkyl, aralkyl, hydroxyalkyl, carboxy or alkoxycarbonyl, or R¹⁰ and R¹¹ show a group which forms cycloalkyl in combination and l, m and n are each 0 or an integer of 1-3,

in the formula (c),

L is hydrogen, alkyl, aminoalkyl, mono- or dialkylaminoalkyl, tetrahydrofurfuryl, carbamoylalkyl,

phthalimidoalkyl, amidino or a group of the formula



wherein B is hydrogen, alkyl, alkoxy, aralkyl, aralkyloxy, aminoalkyl, hydroxyalkyl, alkanoyloxy-alkyl, alkoxycarbonylalkyl, α -aminobenzyl, furyl, pyridyl, phenyl, phenylamino, styryl or imidazopyridyl,

Q^1 is hydrogen, halogen, hydroxy, aralkyloxy or thienylmethyl,

W is alkylene,

Q^2 is hydrogen, halogen, hydroxy or aralkyloxy,

X is alkylene,

Q^3 is hydrogen, halogen, hydroxy, alkoxy, nitro, amino, 2,3-dihydrofuryl or 5-methyl-3-oxo-2,3,4,5-tetrahydropyridazin-6-yl;

and Y is a single bond, alkylene or alkenylene, and

in the formula (c),

a broken line is a single bond or a double bond, and

R^5 is hydrogen, hydroxy, alkoxy, alkoxycarbonyloxy, alkanoyloxy or aralkyloxycarbonyloxy;


Rb is a hydrogen, an alkyl, an aralkyl, an aminoalkyl or a mono- or dialkylaminoalkyl; and

Rc is an optionally substituted heterocycle containing nitrogen,

an isomer thereof and/or a pharmaceutically acceptable acid



addition salt thereof.

3. The agent for the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis of claim 1 or claim 2, wherein the compound having a Rho kinase inhibitory activity is an amide compound of the following formula (I')



wherein

Ra' is a group of the formula




wherein

R' is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring,

R¹ is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring, or R' and R¹ in combination form, together with the adjacent nitrogen atom, a group forming a heterocycle optionally having, in the ring, oxygen atom, sulfur atom or optionally substituted nitrogen atom,

R² is hydrogen or alkyl,

R³ and R⁴ are the same or different and each is hydrogen, alkyl, aralkyl, halogen, nitro, amino, alkylamino, acylamino, hydroxy, alkoxy, aralkyloxy, cyano, acyl, mercapto, alkylthio, aralkylthio, carboxy, alkoxycarbonyl, carbamoyl, alkylcarbamoyl or azide, and
 5 is a group of the formula



wherein R¹⁰ and R¹¹ are the same or different and each is hydrogen, alkyl, haloalkyl, aralkyl, hydroxyalkyl, carboxy or alkoxycarbonyl, or R¹⁰ and R¹¹ show a group which forms cycloalkyl in combination and l, m and n are each 0 or an integer of 1-3,

Rb is a hydrogen, an alkyl, an aralkyl, an aminoalkyl or a mono- or dialkylaminoalkyl; and

Rc is an optionally substituted heterocycle containing
 15 nitrogen,


an isomer thereof and/or a pharmaceutically acceptable acid addition salt thereof.

4. The agent for the prophylaxis and treatment of interstitial
 20 pneumonia and pulmonary fibrosis of claim 1, wherein the compound having a Rho kinase inhibitory activity is a compound selected from the group consisting of (+)-trans-4-(1-aminoethyl)-1-(4-pyridylcarbamoyl)cyclohexane, (+)-trans-N-(1H-pyrrolo[2,3-b]pyridin-4-yl)-4-(1-aminoethyl)cyclohexanecarboxamide, (R)-(+) -
 25 N-(4-pyridyl)-4-(1-aminoethyl)benzamide and (R)-(+) -N-(1H-pyrrolo[2,3-b]pyridin-4-yl)-4-(1-aminoethyl)benzamide, and/or a pharmaceutically acceptable acid addition salt thereof.

5. The agent for the prophylaxis and treatment of interstitial
 30 pneumonia and pulmonary fibrosis of claim 1, wherein the compound




having a Rho kinase inhibitory activity is (+)-trans-4-(1-aminoethyl)-1-(4-pyridylcarbamoyl)cyclohexane and/or a pharmaceutically acceptable acid addition salt thereof.

- 5 6. A pharmaceutical composition for the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis, which comprises a compound having a Rho kinase inhibitory activity and a pharmaceutically acceptable carrier.
- 10 7. The pharmaceutical composition for the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis of claim 6, wherein the compound having a Rho kinase inhibitory activity is an amide compound of the following formula (I)




15 wherein

Ra is a group of the formula




in the formulas (a) and (b),

- 20 R is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring, or a group of the formula



wherein R^6 is hydrogen, alkyl or formula: $-\text{NR}^8\text{R}^9$
 wherein R^8 and R^9 are the same or different and each is
 hydrogen, alkyl, aralkyl or phenyl, R^7 is hydrogen,
 5 alkyl, aralkyl, phenyl, nitro or cyano, or R^6 and R^7 in
 combination show a group forming a heterocycle
 optionally having, in the ring, oxygen atom, sulfur
 atom or optionally substituted nitrogen atom,
 R^1 is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl,
 10 phenyl or aralkyl, which optionally has a substituent
 on the ring, or R and R^1 in combination form, together
 with the adjacent nitrogen atom, a group forming a
 heterocycle optionally having, in the ring, oxygen
 atom, sulfur atom or optionally substituted nitrogen
 15 atom,
 R^2 is hydrogen or alkyl,
 R^3 and R^4 are the same or different and each is hydrogen, alkyl,
 aralkyl, halogen, nitro, amino, alkylamino, acylamino,
 hydroxy, alkoxy, aralkyloxy, cyano, acyl, mercapto,
 20 alkylthio, aralkylthio, carboxy, alkoxycarbonyl,
 carbamoyl, alkylcarbamoyl or azide, and
 A is a group of the formula







wherein R^{10} and R^{11} are the same or different and each is
 25 hydrogen, alkyl, haloalkyl, aralkyl, hydroxyalkyl,
 carboxy or alkoxycarbonyl, or R^{10} and R^{11} show a group

which forms cycloalkyl in combination and l, m and n are each 0 or an integer of 1-3,

in the formula (c),

L is hydrogen, alkyl, aminoalkyl, mono- or dialkylaminoalkyl, tetrahydrofurfuryl, carbamoylalkyl, phthalimidoalkyl, amidino or a group of the formula



wherein B is hydrogen, alkyl, alkoxy, aralkyl, aralkyloxy, aminoalkyl, hydroxyalkyl, alkanoyloxy-alkyl, alkoxycarbonylalkyl, α -aminobenzyl, furyl, pyridyl, phenyl, phenylamino, styryl or imidazopyridyl,

Q^1 is hydrogen, halogen, hydroxy, aralkyloxy or thienylmethyl,

W is alkylene,

Q^2 is hydrogen, halogen, hydroxy or aralkyloxy,

X is alkylene,

Q^3 is hydrogen, halogen, hydroxy, alkoxy, nitro, amino, 2,3-dihydrofuryl or 5-methyl-3-oxo-2,3,4,5-tetrahydropyridazin-6-yl;

and Y is a single bond, alkylene or alkenylene, and

in the formula (c),


a broken line is a single bond or a double bond, and

R^5 is hydrogen, hydroxy, alkoxy, alkoxycarbonyloxy, alkanoyloxy or aralkyloxy carbonyloxy;

Rb is a hydrogen, an alkyl, an aralkyl, an aminoalkyl or



a mono- or dialkylaminoalkyl; and
 Rc is an optionally substituted heterocycle containing
 nitrogen,
 an isomer thereof and/or a pharmaceutically acceptable acid
 5 addition salt thereof.

8. The pharmaceutical composition for the prophylaxis and
 treatment of interstitial pneumonia and pulmonary fibrosis of
 claim 6 or claim 7, wherein the compound having a Rho kinase
 inhibitory activity is an amide compound of the following formula
 (I')



wherein

Ra' is a group of the formula



15

wherein

R' is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl,
 phenyl or aralkyl, which optionally has a substituent
 on the ring,

20 R¹ is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl,
 phenyl or aralkyl, which optionally has a substituent
 on the ring, or R' and R¹ in combination form,

113
5 R²
R³ and R⁴
together with the adjacent nitrogen atom, a group forming a heterocycle optionally having, in the ring, oxygen atom, sulfur atom or optionally substituted nitrogen atom,


is hydrogen or alkyl,

are the same or different and each is hydrogen, alkyl, aralkyl, halogen, nitro, amino, alkylamino, acylamino, hydroxy, alkoxy, aralkyloxy, cyano, acyl, mercapto, alkylthio, aralkylthio, carboxy, alkoxycarbonyl, carbamoyl, alkylcarbamoyl or azide, and

10

A

is a group of the formula



15

wherein R¹⁰ and R¹¹ are the same or different and each is hydrogen, alkyl, haloalkyl, aralkyl, hydroxyalkyl, carboxy or alkoxycarbonyl, or R¹⁰ and R¹¹ show a group which forms cycloalkyl in combination and l, m and n are each 0 or an integer of 1-3,

20 Rb

is a hydrogen, an alkyl, an aralkyl, an aminoalkyl or a mono- or dialkylaminoalkyl; and

Rc

is an optionally substituted heterocycle containing nitrogen,

an isomer thereof and/or a pharmaceutically acceptable acid

25 addition salt thereof.


9. The pharmaceutical composition for the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis of claim 6, wherein the compound having a Rho kinase inhibitory

activity is a compound selected from the group consisting of (+)-trans-4-(1-aminoethyl)-1-(4-pyridylcarbamoyl)cyclohexane, (+)-trans-N-(1H-pyrrolo[2,3-b]pyridin-4-yl)-4-(1-aminoethyl)cyclohexanecarboxamide, (R)-(+)-N-(4-pyridyl)-4-(1-aminoethyl)benzamide and (R)-(+)-N-(1H-pyrrolo[2,3-b]pyridin-4-yl)-4-(1-aminoethyl)benzamide, and/or a pharmaceutically acceptable acid addition salt thereof.

10. The pharmaceutical composition for the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis of claim 6, wherein the compound having a Rho kinase inhibitory activity is (+)-trans-4-(1-aminoethyl)-1-(4-pyridylcarbamoyl)-cyclohexane and/or a pharmaceutically acceptable acid addition salt thereof.




11. A method of the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis, which comprises administering an effective amount of a compound having a Rho kinase inhibitory activity to a patient.

12. The method of the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis of claim 11, wherein the compound having a Rho kinase inhibitory activity is an amide compound of the following formula (I)




wherein

Ra is a group of the formula



in the formulas (a) and (b),

R is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring, or a group of the formula




wherein R⁶ is hydrogen, alkyl or the formula: -NR⁸R⁹ wherein R⁸ and R⁹ are the same or different and each is hydrogen, alkyl, aralkyl or phenyl, R⁷ is hydrogen, alkyl, aralkyl, phenyl, nitro or cyano, or R⁶ and R⁷ in combination show a group forming a heterocycle optionally having, in the ring, oxygen atom, sulfur atom or optionally substituted nitrogen atom,

R¹ is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring, or R and R¹ in combination form, together with the adjacent nitrogen atom, a group forming a heterocycle optionally having, in the ring, oxygen atom, sulfur atom or optionally substituted nitrogen atom,

R² is hydrogen or alkyl,

R^3 and R^4 are the same or different and each is hydrogen, alkyl, aralkyl, halogen, nitro, amino, alkylamino, acylamino, hydroxy, alkoxy, aralkyloxy, cyano, acyl, mercapto, alkylthio, aralkylthio, carboxy, alkoxycarbonyl, carbamoyl, alkylcarbamoyl or azide, and





A is a group of the formula



wherein R^{10} and R^{11} are the same or different and each is hydrogen, alkyl, haloalkyl, aralkyl, hydroxyalkyl, carboxy or alkoxycarbonyl, or R^{10} and R^{11} show a group which forms cycloalkyl in combination and l , m and n are each 0 or an integer of 1-3,

in the formula (c),


L is hydrogen, alkyl, aminoalkyl, mono- or dialkylaminoalkyl, tetrahydrofurfuryl, carbamoylalkyl, phthalimidoalkyl, amidino or a group of the formula



wherein B is hydrogen, alkyl, alkoxy, aralkyl, aralkyloxy, aminoalkyl, hydroxyalkyl, alkanoyloxy-alkyl, alkoxycarbonylalkyl, α -aminobenzyl, furyl, pyridyl, phenyl, phenylamino, styryl or imidazopyridyl, Q^1 is hydrogen, halogen, hydroxy, aralkyloxy or



- thienylmethyl,
W is alkylene,
Q² is hydrogen, halogen, hydroxy or aralkyloxy,
X is alkylene,
5 Q³ is hydrogen, halogen, hydroxy, alkoxy, nitro, amino,
2,3-dihydrofuryl or 5-methyl-3-oxo-2,3,4,5-
tetrahydropyridazin-6-yl;
and Y is a single bond, alkylene or alkenylene, and
in the formula (c),
10 a broken line is a single bond or a double bond, and
R⁵ is hydrogen, hydroxy, alkoxy, alkoxycarbonyloxy,
alkanoyloxy or aralkyloxycarbonyloxy;
Rb is a hydrogen, an alkyl, an aralkyl, an aminoalkyl or
a mono- or dialkylaminoalkyl; and
15 Rc is an optionally substituted heterocycle containing
nitrogen,
an isomer thereof and/or a pharmaceutically acceptable acid
addition salt thereof.

Sub 20
A H
13. The method of the prophylaxis and treatment of interstitial
pneumonia and pulmonary fibrosis of claim 11 or claim 12, wherein
the compound having a Rho kinase inhibitory activity is an amide
compound of the following formula (I')



wherein

- 25 Ra' is a group of the formula



wherein


AY
cat
5
R' is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring,

R¹ is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring, or R' and R¹ in combination form, together with the adjacent nitrogen atom, a group forming a heterocycle optionally having, in the ring, oxygen atom, sulfur atom or optionally substituted nitrogen atom,

R² is hydrogen or alkyl,

R³ and R⁴ are the same or different and each is hydrogen, alkyl, aralkyl, halogen, nitro, amino, alkylamino, acylamino, hydroxy, alkoxy, aralkyloxy, cyano, acyl, mercapto, alkylthio, aralkylthio, carboxy, alkoxycarbonyl, carbamoyl, alkylcarbamoyl or azide, and

A is a group of the formula



wherein R¹⁰ and R¹¹ are the same or different and each is hydrogen, alkyl, haloalkyl, aralkyl, hydroxyalkyl,

carboxy or alkoxycarbonyl, or R^{10} and R^{11} show a group which forms cycloalkyl in combination and l, m and n are each 0 or an integer of 1-3,

Rb is a hydrogen, an alkyl, an aralkyl, an aminoalkyl or a mono- or dialkylaminoalkyl; and

Rc is an optionally substituted heterocycle containing nitrogen,

an isomer thereof and/or a pharmaceutically acceptable acid addition salt thereof.

10

14. The method of the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis of claim 11, wherein the compound having a Rho kinase inhibitory activity is a compound selected from the group consisting of (+)-trans-4-(1-aminoethyl)-1-(4-pyridylcarbamoyl)cyclohexane, (+)-trans-N-(1H-pyrrolo[2,3-b]pyridin-4-yl)-4-(1-aminoethyl)cyclohexanecarboxamide, (R)-(+)-N-(4-pyridyl)-4-(1-aminoethyl)benzamide and (R)-(+)-N-(1H-pyrrolo[2,3-b]pyridin-4-yl)-4-(1-aminoethyl)benzamide, and/or a pharmaceutically acceptable acid addition salt thereof.

20


15. The method of the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis of claim 11, wherein the compound having a Rho kinase inhibitory activity is a (+)-trans-4-(1-aminoethyl)-1-(4-pyridylcarbamoyl)cyclohexane, and/or a

25 pharmaceutically acceptable acid addition salt thereof.

16. Use of a compound having a Rho kinase inhibitory activity for the production of an agent for the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis.




30

17. The use of claim 16, wherein the compound having a Rho kinase inhibitory activity is an amide compound of the following formula (I)




wherein

Ra is a group of the formula



5 in the formulas (a) and (b),

R is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring, or a group of the formula



10 wherein R⁶ is hydrogen, alkyl or formula: -NR⁸R⁹ wherein R⁸ and R⁹ are the same or different and each is hydrogen, alkyl, aralkyl or phenyl, R⁷ is hydrogen, alkyl, aralkyl, phenyl, nitro or cyano, or R⁶ and R⁷ in combination show a group forming a heterocycle optionally having, in the ring, oxygen atom, sulfur atom or optionally substituted nitrogen atom,


15 R¹ is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring, or R and R¹ in combination form, together with the adjacent nitrogen atom, a group forming a

heterocycle optionally having, in the ring, oxygen atom, sulfur atom or optionally substituted nitrogen atom,

R^2 is hydrogen or alkyl,

5 R^3 and R^4 are the same or different and each is hydrogen, alkyl, aralkyl, halogen, nitro, amino, alkylamino, acylamino, hydroxy, alkoxy, aralkyloxy, cyano, acyl, mercapto, alkylthio, aralkylthio, carboxy, alkoxycarbonyl, carbamoyl, alkylcarbamoyl or azide, and

10 A is a group of the formula







15

wherein R^{10} and R^{11} are the same or different and each is hydrogen, alkyl, haloalkyl, aralkyl, hydroxyalkyl, carboxy or alkoxycarbonyl, or R^{10} and R^{11} show a group which forms cycloalkyl in combination and l, m and n are each 0 or an integer of 1-3,

20

in the formula (c),

L is hydrogen, alkyl, aminoalkyl, mono- or dialkylaminoalkyl, tetrahydrofurfuryl, carbamoylalkyl, phthalimidoalkyl, amidino or a group of the formula



25

wherein B is hydrogen, alkyl, alkoxy, aralkyl,

aralkyloxy, aminoalkyl, hydroxyalkyl, alkanoyloxy-
alkyl, alkoxycarbonylalkyl, α -aminobenzyl, furyl,
pyridyl, phenyl, phenylamino, styryl or
imidazopyridyl,

5 Q^1 is hydrogen, halogen, hydroxy, aralkyloxy or
thienylmethyl,

W is alkylene,

Q^2 is hydrogen, halogen, hydroxy or aralkyloxy,

X is alkylene,

10 Q^3 is hydrogen, halogen, hydroxy, alkoxy, nitro, amino,
2,3-dihydrofuryl or 5-methyl-3-oxo-2,3,4,5-
tetrahydropyridazin-6-yl;

and Y is a single bond, alkylene or alkenylene, and
in the formula (c),

15 a broken line is a single bond or a double bond, and


R^5 is hydrogen, hydroxy, alkoxy, alkoxycarbonyloxy,
alkanoyloxy or aralkyloxycarbonyloxy;

Rb is a hydrogen, an alkyl, an aralkyl, an aminoalkyl or
a mono- or dialkylaminoalkyl; and

20 Rc is an optionally substituted heterocycle containing
nitrogen,



an isomer thereof and/or a pharmaceutically acceptable acid
addition salt thereof.

25 18. The use of claim 16 or claim 17, wherein the compound having
a Rho kinase inhibitory activity is an amide compound of the
following formula (I')



30 wherein

Ra' is a group of the formula



AS
ant
wherein


5 R' is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring,

R^1 is hydrogen, alkyl, or cycloalkyl, cycloalkylalkyl, phenyl or aralkyl, which optionally has a substituent on the ring, or R' and R^1 in combination form,
10 together with the adjacent nitrogen atom, a group forming a heterocycle optionally having, in the ring, oxygen atom, sulfur atom or optionally substituted nitrogen atom,

R^2 is hydrogen or alkyl,

15 R^3 and R^4 are the same or different and each is hydrogen, alkyl, aralkyl, halogen, nitro, amino, alkylamino, acylamino, hydroxy, alkoxy, aralkyloxy, cyano, acyl, mercapto, alkylthio, aralkylthio, carboxy, alkoxycarbonyl, carbamoyl, alkylcarbamoyl or azide, and

20 A is a group of the formula



wherein R^{10} and R^{11} are the same or different and each

AS
cut

5 Rb

is hydrogen, alkyl, haloalkyl, aralkyl, hydroxyalkyl, carboxy or alkoxycarbonyl, or R^{10} and R^{11} show a group which forms cycloalkyl in combination and l, m and n are each 0 or an integer of 1-3,

Rc

is a hydrogen, an alkyl, an aralkyl, an aminoalkyl or a mono- or dialkylaminoalkyl; and is an optionally substituted heterocycle containing nitrogen,

an isomer thereof and/or a pharmaceutically acceptable acid

10 addition salt thereof.

19. The use of claim 16, wherein the compound having a Rho kinase inhibitory activity is a compound selected from the group consisting of (+)-trans-4-(1-aminoethyl)-1-(4-

15 pyridylcarbamoyl)cyclohexane, (+)-trans-N-(1H-pyrrolo[2,3-b]pyridin-4-yl)-4-(1-aminoethyl)cyclohexanecarboxamide, (R)-(+)-N-(4-pyridyl)-4-(1-aminoethyl)benzamide and (R)-(+)-N-(1H-pyrrolo[2,3-b]pyridin-4-yl)-4-(1-aminoethyl)benzamide, and/or a pharmaceutically acceptable acid addition salt thereof.

20

20. The use of claim 16, wherein the compound having a Rho kinase inhibitory activity is a (+)-trans-4-(1-aminoethyl)-1-(4-pyridylcarbamoyl)cyclohexane, and/or a pharmaceutically acceptable acid addition salt thereof.

25

21. A commercial package comprising a pharmaceutical composition for the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis of any of claim 6 to claim 10, and a written matter associated therewith, the written matter stating that the
30 pharmaceutical composition can or should be used for the prophylaxis and treatment of interstitial pneumonia and pulmonary fibrosis.